

# Multiple Cranial Nerve Lesions

Last updated: September 5, 2017

Syndrome	Cranial nerves												Associated lesions
	I	II	III	IV	V	VI	VII	VIII	IX	X	XI	XII	
<b>EXTRAPARENCHYMAL</b>													
<b>Foix</b> (superior orbital fissure)			+	+	V <sub>1</sub>	+							
<b>Tolosa-Hunt</b> (lateral wall of cavernous sinus)			+	+	V <sub>1</sub>	+							
<b>Jacod</b> (retro-sphenoid space)	+	+	+	+	+								
<b>Marcus Gunn</b>			+		+								
<b>Gradenigo</b> (apex of petrous bone)					+	+	±	±					
internal auditory meatus							+	+					
pontocerebellar angle					+		+	+	±	±			ataxia
<b>Vernet</b> (jugular foramen)									+	+	+		
<b>Collet-Sicard</b> (retropharyngeal, posterior laterocondylar space)									+	+	+	+	
<b>Villaret</b> (posterior retroparotid space)									+	+	+	+	Horner's syndrome
<b>Tapia</b>									+			+	

Syndrome	Cranial nerves												Associated lesions
	I	II	III	IV	V	VI	VII	VIII	IX	X	XI	XII	
<b>INTRAPARENCHYMAL</b>													
<b>Jackson</b>										+	+	+	
<b>Schmidt</b>										+	+		
<b>Weber</b> (ventral midbrain syndrome)			+				+ <sup>3</sup>						cerebral peduncle (CHP)
<b>Claude</b>			+										red nucleus or dentato-rubro-thalamic tract
<b>Benedikt</b>			+										red nucleus, cerebral peduncle (CHP)
<b>Nothnagel</b>			+										ipsilateral cerebellar ataxia, dizziness, staggering and rolling gait, often nystagmus
central midbrain syndrome			+										red nucleus, subst. nigra, medial lemniscus
<b>Foville</b>						+							CHP
<b>Millard-Gubler</b>							+						CHP
medial medullary syndrome												+	CHP, medial lemniscus
medial pontine syndrome							+						CHP, medial lemniscus, MLF (internuclear ophthalmoplegia), cerebellar connections
<b>Wallenberg</b> (lateral medullary syndrome)								+ <sup>1</sup>		+	+		lateral medullopontine structures: <i>tr. spinothalamicus,</i> <i>tr. reticulospinalis</i> (sympathetic fibers), <i>vestibular connections,</i> <i>inf. cerebellar peduncle</i>
<b>Marie-Foix</b> (lateral inferior pontine syndrome)								+	+				
lateral superior pontine syndrome													
pseudobulbar paralysis								+	+	+		+	
bulbar paralysis								+	+			+	

CHP - contralateral hemiplegia   <sup>1</sup>only nucl. tractus solitarii (taste)   <sup>2</sup>nucl. sensorii of CN5   <sup>3</sup>supranuclear CN7 palsy

INTRAPARENCHYMAL lesions - **crossed** sensory or motor paralysis (cranial nerve signs on one side of body and tract signs on opposite side).

Lesions on BRAINSTEM SURFACE:

- involvement of **adjacent** cranial nerves (often occurring in succession);
- late and rather slight involvement of **long pathways** (sensory and motor) and segmental structures lying within brainstem.

EXTRA-AXIAL lesions - likely to cause **bone erosion** (e.g. enlargement of foramen of exit of cranial nerves); causes of multiple extra-axial cranial nerves involvement:

- 1) diabetes
- 2) trauma
- 3) tumors
- 4) localized infections (e.g. herpes zoster)
- 5) granulomatous disease (e.g. Wegener's granulomatosis), Behçet's disease, sarcoidosis, chronic glandular tuberculosis (scrofula)
- 6) enlarging saccular aneurysms
- 7) platybasia, basilar skull invagination, Chiari malformation.

IDIOPATHIC MULTIPLE CRANIAL NERVE INVOLVEMENT

- on one or both sides of face.
- subacute onset of boring facial **pain** → **paralysis** of motor cranial nerves.
- clinical features overlap those of *Tolosa-Hunt syndrome*.
- frequently responsive to **steroids**.

BIBLIOGRAPHY for ch. "Cranial Neuropathies" → follow this [LINK >>](#)